8. Kemppainen T, Kroger H, Janatuinene E, et al. Bone recovery after a gluten-free diet: a 5-year follow-up study. *Bone*. 1999;25:355-360.
9. Nuti R, Martini G, Valenti R, Giovani S, Salvadori S, Avanzati A. Prevalence of undiagnosed coeliac syndrome in osteoporotic women. *J Intern Med*. 2001; 250:361-366.
10. Mather KJ, Meddings JB, Beck PL, Scott RB, Hanley DA. Prevalence of IgA-antiendomysial antibody in asymptomatic low bone mineral density. *Am J Gastroenterol*. 2001;96:120-125.
11. Lindh E, Ljunghall S, Larsson K, Lavo B. Screening for antibodies against gliadin in patients with osteoporosis. *J Intern Med*. 1992;231:403-406.
12. Gonzalez D, Sugai E, Gomez JC, et al. Is it necessary to screen for celiac disease in postmenopausal osteoporotic women? *Calcif Tissue Int*. 2002;71:141-144.
13. WHO Study Group on Assessment of Fracture Risk and Its Application to Screening for Postmenopausal Osteoporosis. *Assessment of Fracture Risk and Its Application to Screening for Postmenopausal Osteoporosis: Report of a WHO Study Group*. Geneva, Switzerland: World Health Organization; 1994. WHO Technical Report Series 843.
14. Grossman DA, Corazziari E, Tally NJ, Thompson WG, Whitehead WE, eds. *Rome II: The Functional Gastrointestinal Disorders*. 2nd ed. McLean, Va: Degnon Associates Inc; 2000:351-396.
15. Selby PL, Davies M, Adams JE, Mawer EB. Bone loss in celiac disease is related to secondary hyperparathyroidism. *J Bone Miner Res*. 1999;14:652-657.
16. Fabiani E, Catassi C; International Working Group on Eu-tTG. The serum IgA class anti-tissue transglutaminase antibodies in the diagnosis and follow up of coeliac disease: results of an international multi-centre study. *Eur J Gastroenterol Hepatol*. 2001;13:659-665.
17. Korponay-Szabo IR, Dahlbom I, Laurila K, et al. Elevation of IgG antibodies against tissue transglutaminase as a diagnostic tool for coeliac disease in selective IgA deficiency. *Gut*. 2003;52:1567-1571.
18. Looker AC, Orwell ES, Johnston CC Jr, et al. Prevalence of low femoral bone density in older U.S. adults from NHANES III. *J Bone Miner Res*. 1997;12:1761-1768.
19. Fasano A, Berti I, Gerarduzzi T, et al. Prevalence of celiac disease in at-risk and not-at-risk groups in the United States: a large multicenter study. *Arch Intern Med*. 2003;163:286-292.
20. West J, Logan RFA, Hill PG, et al. Seroprevalence, correlates, and characteristics of undetected coeliac disease in England. *Gut*. 2003;52:960-965.

Correction

Duplicate Publication: The letter to the editor by Alexander C. Chester, MD, titled "A Single-Payer National Health Insurance: We Gave Twice at the Office" was inadvertently published twice in the August 9/23 (2004; 164:1702) and November 8 (2004;164:2281) issues of the ARCHIVES.

©2005 American Medical Association. All rights reserved.

Downloaded From: http://archinte.jamanetwork.com/ on 04/01/2015