

clude the following: eliminate mandates for direct physician note entry; provide alternatives, such as dictation, document scanning,⁹ and/or scribes¹⁰ for EMR note capture; make the EMR inbox smarter about when, and how, to present messages to PCPs; and train clinic support personnel to manage the same proportion of inbox messages they handled with paper systems and to think whether a message is important or just noise before pushing the so-convenient "Forward-to-PCP" button. If changes are not made to reduce or eliminate these time penalties on PCPs, there will be no PCPs left to penalize.

Clement J. McDonald, MD
Michael H. McDonald, MD

Author Affiliations: Lister Hill National Center for Biomedical Communications, National Library of Medicine, National Institutes of Health, Bethesda, Maryland (Dr C. J. McDonald); and Department of Surgery, University of Wisconsin School of Medicine (Dr M. H. McDonald).

Correspondence: Dr M. H. McDonald, 3209 Dryden Dr, Madison, WI 53704 (mmcdonal@wisc.edu).

Financial Disclosure: None reported.

Funding/Support: This work was supported in part by the Intramural Research Program of the National Institutes of Health, National Library of Medicine.

Disclaimer: The findings and conclusions in this Invited Commentary are those of the authors and do not necessarily represent the official positions of National Library of Medicine, the National Institutes of Health, the Department of Health and Human Services, or the University of Wisconsin School of Medicine.

Additional Information: Parties interested in obtaining the full survey instrument should contact the corresponding author.

Additional Contributions: Matthew Hoffman, an MD and PhD student at the University of Wisconsin School of Medicine, organized the institutional review board exemption request at the University of Wisconsin and provided suggestions for improving the manuscript.

1. McDonald CJ. Protocol-based computer reminders, the quality of care and the non-perfectability of man. *N Engl J Med*. 1976;295(24):1351-1355.
2. Tierney WM, Miller ME, Overhage JM, McDonald CJ. Physician inpatient order writing on microcomputer workstations. Effects on resource utilization. *JAMA*. 1993;269(3):379-383.
3. Murphy DR, Reis B, Kadiyala H, et al. Electronic health record-based messages to primary care providers: valuable information or just noise? *Arch Intern Med*. 2012;172(3):283-285.
4. Murphy DR, Reis B, Sittig DF, Singh H. Notifications received by primary care practitioners in electronic health records a taxonomy and time analysis. *Am J Med*. 2012;125(2):209.e1-209.e7.
5. R Development Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing; 2010.
6. Chaudhry B, Wang J, Wu S, et al. Systematic review: impact of health information technology on quality, efficiency, and costs of medical care. *Ann Intern Med*. 2006;144(10):742-752.
7. Pizziferri L, Kitzler AF, Volk LA, et al. Primary care physician time utilization before and after implementation of an electronic health record: a time-motion study. *J Biomed Inform*. 2005;38(3):176-188.
8. Hauer KE, Durning SJ, Kernan WN, et al. Factors associated with medical students' career choices regarding internal medicine. *JAMA*. 2008;300(10):1154-1164.
9. Downs SM, Biondich PG, Anand V, Zore M, Carroll AE. Using Arden Syntax and adoptive turnaround documents to evaluate clinical guidelines. *AMIA Annu Symp Proc*. 2006:214-218.
10. Conn J. Getting it in writing: docs using scribes to ease the transition to EHRs. *Mod Healthc*. 2010;40(6):30-32.

RESEARCH LETTERS

ONLINE FIRST

Young-Onset Colorectal Cancer: Is It Time to Pay Attention?

The 2010 *Annual Report to the Nation on Cancer* celebrated a steady decline in the incidence of colorectal cancer (CRC).¹ This progress has been largely attributed to CRC screening, recommended for adults 50 years or older since 1996.² In sharp contrast to overall trends, the incidence of CRC appears to be increasing among adults younger than 50 years,^{1,3} a group for whom average-risk screening is not routine. Particularly concerning is a trend toward advanced-stage CRCs,^{1,3} suggesting a potential role for increased clinical vigilance and more prompt evaluation of symptomatic patients. To raise clinical awareness and facilitate recognition, we undertook a cohort study to (1) examine incidence trends; (2) define the distinct clinicopathologic manifestations of young-onset CRC; and (3) identify risk factors for advanced-stage disease.

Methods. We used the National Cancer Database (NCDB), a hospital-based cancer registry sponsored by the American College of Surgeons and American Cancer society that captures 70% of all incident cancers annually.⁴ We identified patients diagnosed as having invasive adenocarcinoma (behavior code 3 [malignant]; histology codes 8010, 8020-8022, 8140-8144, 8210-8211, 8260-8263, 8470-8473, 8480-8481, and 8490) of the colon and the rectum between January 1998 and December 2007, the most recent decade after the US Preventive Services Task Force (USPSTF) recommended CRC screening. Eligible patients (n=588 869) were stratified by tumor site (colon [site codes C180, 182-189, and 199] vs rectum [site code C209]) and by age at diagnosis (young onset [before age 50 years] vs later onset [at age 50 years or older]). Age-adjusted incidence rates (per 100 000 individuals) were calculated using the 2000 US standard population (SEER*Stat software, version 7.0.4; National Cancer Institute). The temporal trend in the annual percentage change (APC) was identified (Joinpoint Regression Program, version 3.5.1; National Cancer Institute). For a subgroup (n=34 781) of young-onset CRCs diagnosed between 2003 and 2007—the following years after the USPSTF strengthened its screening recommendation in 2002⁵—independent predictors of advanced-stage disease (stages III and IV, nodal and distant metastatic disease) were identified by multiple logistic regression. This study was granted exemption status by The University of Texas MD Anderson Cancer Center institutional review board.

Results. We identified 64 068 young-onset (10.9%) and 524 801 later-onset (89.1%) CRCs. Age-adjusted incidence declined for later-onset CRCs after 2001 (APC, -2.5%; 95% CI, -3.0% to -2.0%) but has consistently increased since 2001 for young-onset disease (APC, 2.1%; 95% CI, 1.1% to 3.1%). The increase was greater for

young-onset rectal (APC, 3.9%; 95% CI, 3.1% to 4.7%) than colon (APC, 2.7%; 95% CI, 2.0% to 3.3%) cancers. The median age for young-onset CRCs was 44 years, with most (75.2%) occurring between ages 40 and 49 years. Compared with later-onset disease, young-onset CRCs were more prevalent among patients with nonwhite race/ethnicity (29.5% vs 17.6%; $P < .001$), who were not insured or insured by Medicaid (16.5% vs 4.7%; $P < .001$) and who lived in the southern and western parts of the United States (56.2% vs 50.3%; $P < .001$). Young-onset CRC more commonly arose from colon distal to the splenic flexure or the rectum (69.0% vs 57.7%; $P < .001$). Mucinous and signet-ring histologic subtype (12.6% vs 10.8%; $P < .001$) and poor or no differentiation (20.4% vs 18%; $P < .001$) were also more frequently exhibited. Advanced-stage disease was diagnosed significantly more commonly in young patients (Table). Independent risk factors for advanced-stage disease included younger age (age 30-39 years: hazard ratio [HR], 1.21; 95% CI, 1.1 to 1.4; age 18-29 years: HR, 1.4; 95% CI, 1.2 to 1.6; vs age 40-49 years); African American race (HR, 1.2; 95% CI, 1.1 to 1.3; vs white race) and lack of insurance or Medicaid insurance (uninsured: HR, 1.2; 95% CI, 1.1 to 1.3; Medicaid: HR, 1.6; 95% CI, 1.5 to 1.8; vs insured).

Comment. This cohort study of young-onset CRC, the largest to date, highlights an alarming rise in incidence since 2001. Most worrisome is the high proportion of stage III/IV disease (Figure), consistent with a prior smaller population-based report.⁶ In the absence of routine screening, contributing factors to these trends may include (1) a reluctance on the part of young adults to seek medical care⁷; (2) the large percentage of young adults without insurance or ready access to care (18.3%-29.2% for adults aged 18-44 years vs 1.3%-13.6% for adults older than 45 years⁸); and (3) an underappreciation of the increasing risk for young-onset CRC, leading clinicians to overlook or dismiss symptoms that are nonspecific but may be consistent with CRC (ie, rectal bleeding, abdominal pain or cramping, change in bowel pattern). Finally, the predilection of young-onset CRCs for the distal colon and rectum identify these as high-yield anatomic regions for endoscopic evaluation in symptomatic pa-

tients and as potentially cost-effective targets for screening programs in presymptomatic young adults.

This study could not establish the mechanisms underlying observed sociodemographic disparities, nor did it explore the molecular basis of young-onset CRC (familial adenomatous polyposis was excluded). Notwithstanding, these data argue for heightened awareness of these concerning trends in young-onset CRC. Symptomatic young patients should undergo timely sigmoidoscopy at a minimum, if not a full colonoscopy. Identifying high-risk cohorts for targeted screening should be a priority.

Y. Nancy You, MD, MHS
 Yan Xing, MD, PhD
 Barry W. Feig, MD
 George J. Chang, MD, MS
 Janice N. Cormier, MD, MPH

Published Online: December 12, 2011. doi:10.1001/archinternmed.2011.602. This article was corrected for an error in the Figure key on December 13, 2011.

Author Affiliations: Department of Surgical Oncology, The University of Texas MD Anderson Cancer Center, Houston.

Correspondence: Dr You, Department of Surgical Oncology, Unit 1484, University of Texas MD Anderson Cancer Center, 1400 Pressler St, PO Box 301402, Houston, TX 77230-1402 (ynyou@mdanderson.org).

Author Contributions: Dr You had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of data analysis. *Study concept and design:* You, Xing, Chang, and Cormier. *Acquisition of data:* You, Xing, and Cormier. *Analysis and interpretation of data:* You, Xing, Feig, Chang, and Cormier. *Drafting of the manuscript:* You and Cormier. *Critical revision of the manuscript for important intellectual content:* You, Xing, Feig, Chang, and Cormier. *Statistical analysis:* You, Xing, and Cormier. *Obtained funding:* You and Cormier. *Study supervision:* Feig and Cormier. **Financial Disclosure:** Dr Feig served on the speakers bureau for and received honorarium from Genomic Health. **Funding/Support:** This study was supported by the University of Texas MD Anderson Cancer Center Institutional Research Grant (Dr You) and The University of Texas MD Anderson Cancer Center's Institute for Cancer Care Excellence.

Previous Presentation: This study was presented in part at the Sixth Annual Academic Surgical Congress, Association for Academic Surgery; February 4, 2011; Orange County, California.

Additional Contributions: Miguel A. Rodriguez-Bigas, MD, and John M. Skibber, MD, assisted with critical revision of the manuscript, and Yan K. Ling, MPH, assisted with data analysis.




Figure. Distribution of cancer stage at diagnosis for young-onset vs later-onset colon and rectal cancers.

1. Edwards BK, Ward E, Kohler BA, et al. Annual report to the nation on the status of cancer, 1975-2006, featuring colorectal cancer trends and impact of interventions (risk factors, screening, and treatment) to reduce future rates. *Cancer*. 2010;116(3):544-573.
2. US Preventive Services Task Force. *Guide to Clinical Preventive Services: Report of the US Preventive Services Task Force*. 2nd ed. Washington, DC: Dept of Health and Human Services; 1995.
3. Siegel RL, Jemal A, Ward EM. Increase in incidence of colorectal cancer among young men and women in the United States. *Cancer Epidemiol Biomarkers Prev*. 2009;18(6):1695-1698.
4. Winchester DP, Stewart AK, Phillips JL, Ward EE. The national cancer data base: past, present, and future. *Ann Surg Oncol*. 2010;17(1):4-7.

- US Preventive Services Task Force. Screening for colorectal cancer: recommendation and rationale. *Ann Intern Med.* 2002;137(2):129-131.
- O'Connell JB, Maggard MA, Liu JH, Etzioni DA, Livingston EH, Ko CY. Rates of colon and rectal cancers are increasing in young adults. *Am Surg.* 2003;69(10):866-872.
- Bleyer A. CAUTION! Consider cancer: common symptoms and signs for early detection of cancer in young adults. *Semin Oncol.* 2009;36(3):207-212.
- US Census Bureau. *Income, Poverty, and Health Insurance Coverage in the United States.* 2006. <http://www.census.gov/hhes/www/income/income.html>. Accessed April 31, 2011.

ONLINE FIRST

Supplemental Oxygen Therapy in Medical Emergencies: More Harm Than Benefit?

In medical emergencies, such as acute coronary syndrome, cardiopulmonary resuscitation (CPR), stroke, and exacerbations of chronic obstructive pulmonary disease (COPD), supplemental oxygen is often routinely administered. Most physicians believe this intervention is potentially lifesaving, and many guidelines support the routine use of high-dose supplemental oxygen.

Over the decades, however, potential detrimental effects of supplemental oxygen appear to have been ignored. Many clinicians are unaware of the variety of pre-clinical studies that have been executed, showing that hyperoxia causes both coronary and systemic vasoconstriction, resulting in deterioration of several important (hemodynamic) parameters (**Table**). The prime candidate mechanism for these unintended effects is believed to be the formation of reactive oxygen species. In this Research Letter, we draw attention to the collective clinical evidence, which argues against the routine use of high-

dose oxygen. Awaiting more thorough studies, we strongly recommend a policy of careful, titrated oxygen supplementation.

Methods. We conducted a search of the literature in MEDLINE and EMBASE to identify articles addressing the effect of oxygen therapy in acute coronary syndrome, cardiopulmonary resuscitation, stroke, and exacerbations of chronic obstructive pulmonary disease.

Results. Large (randomized) clinical studies addressing oxygen supplementation are scarce. In 1976, a double-blind randomized trial was performed in 200 patients with suspected acute myocardial infarction. In the supplemental oxygen group, 9 of 80 patients (11%) died, as opposed to 3 of 77 (3.9%) in patients breathing compressed air (relative risk [RR] of mortality, 2.9; 95% CI, 0.8-10.3). A recent Cochrane review combined this trial with a smaller similar one, which generated a composite RR of mortality of 3.03 (95% CI, 0.93-9.83).¹ In acute decompensated heart failure, no clinical studies are available, despite the rather abundant evidence from preclinical studies, suggesting that such patients may experience the adverse effects caused by coronary and systemic vasoconstriction (Table).

In accordance with existing guidelines, supplemental oxygen is often administered during CPR. In the postresuscitation phase, evidence exists that 30% oxygen is more brain protective than pure oxygen.² Recently, an observational study in 6326 patients showed that supplemental oxygen induced postresuscitation hyperoxia was independently associated with increased mortality (odds ratio [OR], 1.8; 95% CI, 1.5-2.2). A subsequent analysis of the same cohort indicated that each 25-mm Hg increase in PaO₂ was associated with a sta-

Table. Effects of Supplemental Oxygen in Different Clinical Conditions

Clinical Condition	Effect Parameter	Effect	Reference
Exercise testing	ECG alterations	Prolonged	<i>JAMA.</i> 1950;144:373-375
	Heart rate	Decreased	<i>Lancet.</i> 1964;2(7364):825-832
Post myocardial infarction	Stroke volume	Decreased	<i>Br Heart J.</i> 1965;27:401-407
	Cardiac output	Decreased	<i>Br Med J.</i> 1968;4(5627):360-364
	Systemic vascular resistance	Increased	
	Mean arterial pressure	Increased	
Acute myocardial infarction	Vascular resistance of LAD	Increased	<i>J Appl Physiol.</i> 2007;102(5):2040-2045
	Coronary vascular resistance	Increased	<i>Am J Physiol.</i> 2005;288(3):H1057-H1062
Cardiac catheterization	Coronary blood flow	Decreased	
	Stroke volume	Decreased	<i>J Am Coll Cardiol.</i> 1996;27(2):353-357
Congestive heart failure	Heart rate	Decreased	<i>Am J Physiol Heart Circ Physiol.</i> 2002;282(6):H2414-2421
	Cardiac output	Decreased	<i>Chest.</i> 2001;120(2):467-473
	Systemic vascular resistance	Increased	<i>Heart.</i> 2010;96(7):533-538
	LV end diastolic pressure	Increased	
	Isovolumetric relaxation time	Increased	
	COPD	Hypercapnia	Increased
Stroke	Mortality	Increased	<i>BMJ.</i> 2010;341:c5462
	Stroke severity score	Increased	<i>Stroke.</i> 2003;34(2):571-574
	Mortality	Increased	<i>Stroke.</i> 1999;30(10):2033-2037 NCT00414726 ^a
Cardiopulmonary resuscitation	Neuron-specific enolase	Elevated	<i>Resuscitation.</i> 2006;69(2):199-206
	Mortality	Increased	<i>JAMA.</i> 2010;303(21):2165-2171 <i>Circulation.</i> 2011;123(23):2717-2722 <i>Critical Care.</i> 2011;15(2):R90

Abbreviations: COPD, chronic obstructive pulmonary disease; ECG, electrocardiogram; LAD, left anterior descending coronary artery; LV, left ventricular.
^aTrial was terminated (Clinical Trial of Normobaric Oxygen Therapy in Acute Ischemic Stroke, not published, clinicaltrials.gov Identifier: NCT00414726).