HEALTH CARE REFORM Potentially Avoidable 30-Day Hospital Readmissions in Medical Patients

Derivation and Validation of a Prediction Model

Jacques Donzé, MD, MSc; Drahomir Aujesky, MD, MSc; Deborah Williams, MHA; Jeffrey L. Schnipper, MD, MPH

Importance: Because effective interventions to reduce hospital readmissions are often expensive to implement, a score to predict potentially avoidable readmissions may help target the patients most likely to benefit.

Objective: To derive and internally validate a prediction model for potentially avoidable 30-day hospital readmissions in medical patients using administrative and clinical data readily available prior to discharge.

Design: Retrospective cohort study.

Setting: Academic medical center in Boston, Massachusetts.

Participants: All patient discharges from any medical services between July 1, 2009, and June 30, 2010.

Main Outcome Measures: Potentially avoidable 30day readmissions to 3 hospitals of the Partners Health-Care network were identified using a validated computerized algorithm based on administrative data (SQLape). A simple score was developed using multivariable logistic regression, with two-thirds of the sample randomly selected as the derivation cohort and one-third as the validation cohort. **Results:** Among 10731 eligible discharges, 2398 discharges (22.3%) were followed by a 30-day readmission, of which 879 (8.5% of all discharges) were identified as potentially avoidable. The prediction score identified 7 independent factors, referred to as the HOSPITAL score: *h*emoglobin at discharge, discharge from an *oncology* service, sodium level at discharge, *p*rocedure during the index admission, index *type* of admission, number of *a*dmissions during the last 12 months, and length of stay. In the validation set, 26.7% of the patients were classified as high risk, with an estimated potentially avoidable readmission risk of 18.0% (observed, 18.2%). The HOSPITAL score had fair discriminatory power (*C* statistic, 0.71) and had good calibration.

Conclusions and Relevance: This simple prediction model identifies before discharge the risk of potentially avoidable 30-day readmission in medical patients. This score has potential to easily identify patients who may need more intensive transitional care interventions.

JAMA Intern Med. 2013;173(8):632-638. Published online March 25, 2013. doi:10.1001/jamainternmed.2013.3023



OSPITAL READMISSIONS ARE common and costly. Recent studies estimated a 30-day readmission rate in the United States of 18%;

among Medicare beneficiaries, readmissions are estimated to cost \$17 billion annually.¹ Because at least some hospital readmissions may be avoidable, readmission rates are now used for benchmarking across hospitals, with financial penalties for hospitals with high risk-adjusted rates.²

Several interventions have been shown³⁻⁶ to be effective in reducing the rate of readmission. To improve efficiency, the highest intensity interventions should be targeted to patients who are most likely to benefit. Few models exist to predict 30-day readmission risk in general medical patients.⁷⁻¹⁰ These models do not distinguish between avoidable and unavoidable readmissions, often have poor discriminatory power or calibration, and/or use complex scores not calculable before hospital discharge.



See also pages 624 and 629

To help clinicians target transitional care interventions most efficiently, we derived and internally validated a prediction model for potentially avoidable 30-

JAMA INTERN MED/VOL 173 (NO. 8), APR 22, 2013 WWW.JAMAINTERNALMED.COM

Author Affiliations: Division of

General Medicine and Primary

Care (Drs Donzé and Schnipper

(Dr Schnipper), Brigham and

Massachusetts; Harvard Medical

School, Boston (Drs Donzé and

General Internal Medicine, Bern

Women's Hospital, Boston,

Schnipper); and Division of

University Hospital, Bern,

Switzerland (Dr Aujesky).

and Ms Williams) and

Hospitalist Service

day hospital readmissions in medical patients using readily available administrative and clinical data.

METHODS

STUDY DESIGN AND POPULATION

This retrospective cohort study included consecutive adult patient discharges from all medical services of the Brigham and Women's Hospital with a discharge date between July 1, 2009, and June 30, 2010. Brigham and Women's Hospital is a 750-bed academic medical center in Boston, Massachusetts, with 44 000 inpatient admissions per year. Medical services include general medicine, cardiology, oncology, bone marrow transplant, endocrinology, gastroenterology, hematology, infectious diseases, rheumatology, and nephrology. To avoid inclusion of observation stays, only hospitalizations with a length of stay of more than 24 hours were included. We also excluded hospitalizations when the patient died before discharge, was transferred to another acute health care facility, or left against medical advice. The hospitalizations were randomized into a derivation set (two-thirds of admissions) and a validation set (one-third of admissions). The protocol was approved by the institutional review board of Brigham and Women's Hospital/Partners HealthCare.

STUDY OUTCOME

Among the included patients, there were 3 possible outcomes: admissions not followed by any 30-day readmission, admissions followed by a 30-day potentially avoidable readmission, and admissions followed by a 30-day unavoidable readmission. Because we were interested in the identification of predictors specific to avoidable readmissions, and to give a clear contrast, we chose to compare the admissions followed by a potentially avoidable readmission with those not followed by any 30-day readmission. We excluded the unavoidable readmissions because they correspond to an intermediate and heterogeneous population of patients (eg, some of these patients might have subsequently developed an avoidable readmission had they not already been readmitted). Hospital readmissions were considered to be unavoidable if any of the following characteristics were present: (1) planned readmission (eg, scheduled at the time of the index admission, planned treatment follow-up, and planned chemotherapy) or (2) unforeseen readmission for newly developed conditions not related to known diseases during the index hospitalization.

To exclude the unavoidable readmissions meeting these criteria, we used a validated computerized algorithm (Striving for Quality Level and Analyzing of Patient Expenses [SQLape], developed by Yves Eggli, MD) commonly used in Switzerland to benchmark and compare hospitals.¹¹ This algorithm is based on administrative data and International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes of both the index admission and readmission to identify unavoidable readmissions as described in the previous paragraph. Conversely, the algorithm identifies as potentially avoidable readmissions related to a previously coded medical condition or resulting from a complication of treatment (eg, deep vein thrombosis). The sensitivity as well as specificity of the screening algorithm reached 96% when compared with medical record review (using the same criteria) in a random sample of admission-readmission pairs drawn from the university hospital from which the tool was originally derived.12 Using medical record review, the original SQLape investigators estimated that 23% of potentially avoidable readmissions (1.2% of all readmissions) were clearly avoidable had different action been taken in the hospital (eg, premature discharge, erroneous diagnosis, inappropriate treatment, or inadequate patient education).

Because administrative data may not always accurately identify elective from nonelective readmissions, all potentially avoidable readmissions identified with SQLape were further reviewed by 1 of 9 trained senior medical residents from Brigham and Women's Hospital to exclude patients with a planned readmission.

Because potentially avoidable readmissions to departments and hospitals other than the Department of Medicine of Brigham and Women's Hospital could occur, we looked at readmissions to any service of Brigham and Women's Hospital, the Massachusetts General Hospital, or the Faulkner Hospital, all of which are affiliated with the Partners HealthCare network. Each admission could be considered as both an index admission and a readmission when several readmissions occurred within an interval of less than 30 days.

PREDICTOR VARIABLES

We collected data on several types of variables from easily obtainable sources (**Table 1**), including demographic information, previous health care utilization, primary care provider information, and index admission characteristics from administrative data sources; procedures and chronic medical conditions from billing data; and last known laboratory values before discharge. Variables were chosen a priori and according to the medical literature.^{8,13,14}

STATISTICAL ANALYSIS

The presence of any difference in baseline characteristics between the groups with a 30-day potentially avoidable readmission and those not readmitted was tested by univariable logistic regression. Using the derivation set, all variables noted in Table 1 were included in an initial multivariable model. When laboratory values were missing (<1.1% of discharges), these variables were considered as within normal limits. Because the same patient could have had several admissions and readmissions during the study period, data were clustered at the patient level using general estimating equations. The less strongly linked variables were removed from the model one at a time by backward elimination until all predictors were significant at P < .05.

The result of the multivariable regression model was then used to develop a prediction score by using a regression coefficient–based scoring method.¹⁵ Integer scores were assigned by dividing risk-factor coefficients by the smallest coefficient and rounded up to the nearest integer. After model derivation, the risk for an admission to be followed by a 30-day potentially avoidable readmission was categorized into 3 groups (low, intermediate, and high) for ease of interpretation.

The discriminatory power of the resulting score was assessed in both the derivation and validation set by calculating the cross-validated C statistic,16 which refers to the ability to differentiate between admissions followed and those not followed by a 30-day potentially avoidable readmission. How closely the predicted probabilities reflected actual risk was assessed by calibration and with the Hosmer-Lemeshow goodnessof-fit test (nonsignificant P values on this test imply good fit). Details of calibration are shown by comparing the estimated risk obtained with the score for patients in a specific category with the observed probability, ie, the actual proportion of 30day potentially avoidable readmissions for patients in that category. Finally, because in practice unavoidable readmissions cannot be identified and excluded before risk estimation, we reevaluated the performance of the score when applied to the complete cohort, including both potentially avoidable and unavoidable readmissions. Analyses were performed with commercial software (SAS, version 9.2; SAS Institute, Inc).

Table 1. Baseline Characteristics of the Entire Cohort and Univariable Analysis in the Derivation Set

	No. (%)			
		Univariable Analyses in the Derivation Set (n = 6141)		
Characteristic	Entire Cohort (N = 9212)	No Readmission (n = 5553)	PAR (n = 588)	P Value
Age >75 y ^a	2051 (22.3)	1277 (23.0)	119 (20.2)	.16
Male sex	4476 (48.6)	2652 (47.8)	282 (48.0)	.93
Ethnicity	CCEE (70.0)	2020 (71.0)		
Non-Hispanic black	1498 (16 3)	011 (16 A)	427 (72.0)	
Hispanic	773 (8.4)	467 (8.4)	53 (9.0)	.32
Other	286 (3.1)	186 (3.3)	9 (1.5)	
First language				
English ^a	8376 (90.9)	5043 (90.8)	536 (91.2)	10
Spanish	482 (5.2)	285 (5.1)	37 (6.3)	.42
Utiter Marital status	354 (3.8)	225 (4.1)	15 (2.5)	
Current spouse or partner ^a	4695 (51.0)	2792 (50.3)	305 (51.9)	
Single/never married	2315 (25.1)	1385 (24.9)	149 (25.3)	.36
Separated/divorced/widowed/no answer	2202 (23.9)	1376 (24.8)	134 (22.8)	
Primary insurance	. ,		. ,	
Medicare ^a	4676 (50.8)	2811 (50.6)	281 (47.8)	
Medicaid	749 (8.1)	443 (8.0)	55 (9.4)	.31
Private	3775 (41.0)	2291 (41.3)	249 (42.3)	101
NORe Source of index admission	12 (0.1)	8 (0.1)	3 (0.5)	
Emergency department ^a	4902 (53.2)	2938 (52 9)	315 (53.6)	
Direct from home/outpatient clinic	2830 (30 7)	1719 (31.0)	183 (31 1)	73
Nursing home/rehabilitation facility/other hospital	1480 (16.1)	896 (16.1)	90 (15.3)	
Type of index admission				
Elective ^a	1188 (12.9)	736 (13.3)	64 (10.9)	11
Nonelective	8021 (87.1)	4817 (86.7)	524 (89.1)	.11
Division of index admission	0100 (00.0)		000 (00 5) -	
Oncology	2192 (23.8)	1204 (21.7)	232 (39.5)	<.001
Uther medical service"	7020 (76.2)	4349 (78.3)	356 (60.5)	
1-4 ^a	5181 (56.2)	3252 (58.6)	259 (44 0)	
>4	4031 (43.8)	2301 (41.4)	329 (56.0)	<.001
No. of hospital admissions in the past year				
0 ^a	4321 (46.9)	2698 (48.6)	178 (30.3) 🗍	
1-5	4456 (48.4)	2629 (47.3)	344 (58.5)	<.001
>5	435 (4.7)	226 (4.1)	66 (11.2)	
Identified caregiver at discharge	8459 (91.8)	5089 (91.6)	551 (93.7)	.10
No. of medications at discharge	1863 (20.2)	118/ (21.3)	80 (15 1) ¬	
7-9	2017 (21.9)	1233 (22.2)	109 (18 5)	
10-13	2503 (27.2)	1482 (26.7)	152 (25.9)	<.001
>13	2829 (30.7)	1654 (29.8)	238 (40.5)	
No. of procedures during index admission		. ,	. ,	
0 ^a	3636 (39.5)	2263 (40.8)	177 (30.1)	< 001
≥1	5576 (60.5)	3290 (59.2)	411 (69.9)	
Hemoglobin level at discharge, g/dL ^b		0701 (07 7)		
<12.0 >10.0a	5626 (61.1)	3/61 (67.7)	481 (81.8)	<.001
\leq 12.0 ⁻⁴ Serum sodium level at discharge mEg/l ^c	5560 (56.9)	1792 (32.3)	107 (10.2)	
	1454 (15.8)	832 (15.0)	137 (23.3)	
≥135 ^a	7758 (84.2)	4721 (85.0)	451 (76.7)	<.001
GFR at discharge, mL/min ^d	()	()		
<30	892 (9.7)	527 (9.5)	76 (12.9)	
30-59	1977 (21.5)	1178 (21.2)	119 (20.2)	.07
$\geq 60^{a}$	6343 (68.8)	3848 (69.3)	393 (66.8)	
Comorbidity	0010 (05.1)	1070 (04.0)	164 (07.0)	45
Ischemic heart disease [†]	2312 (25.1)	15/9 (24.8)	104 (27.9)	.15
וסטושוווט וולמון עוסלמסל	2497 (27.1)	1300 (27.2)	139 (23.0)	.09

(continued)

JAMA INTERN MED/VOL 173 (NO. 8), APR 22, 2013 WWW.JAMAINTERNALMED.COM

634

634

Table 1. Baseline Characteristics of the Entire Cohort and Univariable Analysis in the Derivation Set (continued)

		No. (%)			
		Univariable Analyses in the Derivation Set (n = 6141)			
Characteristic	Entire Cohort (N = 9212)	No Readmission (n = 5553)	PAR (n = 588)	P Value	
Comorbidity (ct)					
Heart failure ^g	2029 (22.0)	1221 (22.0)	142 (24.1)	.27	
Atrial fibrillation ^h	1633 (17.7)	1000 (18.0)	89 (15.1)	.12	
COPD ⁱ	936 (10.2)	567 (10.2)	61 (10.4)	.91	
Malignant neoplasm ^j	3250 (35.3)	1840 (33.1)	300 (51.0)	<.001	

Abbreviations: COPD, chronic obstructive pulmonary disease; GFR, glomerular filtration rate; PAR, potentially avoidable readmission.

SI conversion factors: To convert hemoglobin to grams per liter, multiply by 10; conversion of serum sodium to millimoles per liter is 1:1.

^aReference group in the multivariable logistic regression.

^bSeventy-eight missing values among the entire cohort (0.8%).

^cSeventy-three missing values among the entire cohort (0.8%).

^dNinety-eight missing values among the entire cohort (1.1%); GFR was calculated using the Modified Diet in Renal Disease method.

^e International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes 249.00 through 250.99 for any diagnosis during the index hospitalization.

^f*ICD-9-CM* codes 410.00 through 414.99 for any diagnosis during the index hospitalization.

^g *ICD-9-CM* codes 428.x, 425.4 through 425.9, 402.01, 402.11, and 398.91 for any diagnosis during the index hospitalization.

^h *ICD-9-CM* codes 427.30 through 427.32 for any diagnosis during the index hospitalization.

ⁱ ICD-9-CM codes 491.00 through 492.99, 493.2, and 496 for any diagnosis during the index hospitalization.

^j ICD-9-CM codes 140.00 through 290.99 for any diagnosis during the index hospitalization.

RESULTS

During the study period, a total of 12 383 patients were discharged from the medical services of the Brigham and Women's Hospital (Figure). Of these patients, 1652 admissions (13.3%) were excluded because of death before discharge, because of transfer to another acute health care facility, or because the patient left against medical advice. Among the 10731 remaining discharges, 2398 (22.3%) were followed by a 30-day readmission, of which 879 readmissions (8.5% of all index discharges, 36.7% of readmissions) were identified as potentially avoidable. The admissions not followed by a 30-day readmission (n = 8333) and the potentially avoidable readmissions (n = 879) were together randomly divided into a derivation set (two-thirds [n = 6141]) and a validation set (one-third [n = 3071]). Overall, 7123 unique patients accounted for all 9212 index discharges.

Baseline characteristics are reported in Table 1. The patients' mean age at inclusion was 61.3 years, and approximately half were male. At least 1 procedure was performed in 60.5% of index admissions, with a mean of 1.8 procedures per admission (**Table 2** includes a list of commonly performed procedures).

The backward multivariable logistic regression analysis identified 8 covariates that were significant independent predictors (eTable 1 reports on the model with all a priori identified predictors; http://www.jamainternalmed .com). To simplify the score, we further excluded the covariate congestive heart failure from the model for 2 reasons: (1) not using any comorbidities would greatly simplify the calculation of the score and, in some cases, the diagnosis might be available only after discharge; (2) the covariate congestive heart failure had one of the higher *P* values (.03) in the multivariable logistic regression. We ended up with a 7-factor predictor score, which we refer to as the HOSPITAL score: *h*emoglobin at discharge, discharge from



Figure. Study flow diagram. SQLape indicates Striving for Quality Level and Analyzing of Patient Expenses.

an *o*ncology service, sodium level at discharge, *p*rocedure during the index admission (any *ICD-9-CM*–coded procedure), index *type* of admission (nonelective vs elective), number of admissions during the past 12 months, and length of stay (**Table 3**; eTable 2 lists the β coefficients and odds ratios for each predictor). Using the score, the risk of potentially avoidable readmission was stratified into 3 categories: low, intermediate, and high. Low-risk patients having 0 to 4 points (49.3% of patients) had a 5.2% estimated risk of potentially avoidable readmission and an

Table 2. Most Frequent ProceduresAmong the Index Admissions

Procedure	Frequency, %
Injection or infusion of cancer chemotherapeutics	8.8
Biopsy and closed biopsy	7.5
Transfusion (eg, platelets, packed red blood cells)	7.1
Gastrointestinal endoscopy	6.5
Venous catheterization, not elsewhere classified	6.1
Percutaneous transluminal coronary angioplasty	5.2
Cardiac catheterization of the left and/or right side of the heart	5.2
Autologous or allogenic stem cell transplant without purging	5.0
Catheter ablation of lesion or tissues of heart	2.8
Implantation or replacement of intracardiac defibrillator or pacemaker	2.8
Continuous positive airway pressure	2.7
Hemodialysis	2.7
Magnetic resonance imaging	2.3
Percutaneous abdominal drainage (ie, paracentesis)	2.1
Other radiotherapeutic procedure	1.8
Thoracentesis	1.5
Lumbar puncture	1.4

observed proportion of 5.4% in the derivation set; highrisk patients having 7 or more points (24.4% of patients) had an 18.3% estimated probability of potentially avoidable readmission and an observed probability of 18.7% (Table 4; eTable 3 lists the risk level for each total point score). The results were very similar in the validation set. The Hosmer-Lemeshow goodness-of-fit statistics were P = .28 and P = .15 in the derivation and validation sets, respectively, indicating good calibration (eTable 4 reports observed and expected results by decile of risk). The discriminatory power of the score was fair, with a crossvalidated C statistic of 0.69 in the derivation set and 0.71 in the validation set. When the HOSPITAL score was applied to the complete cohort before exclusion of unavoidable readmissions (n = 10731), discrimination remained fair, with a cross-validated C statistic of 0.67 and very good calibration (eTable 5).

COMMENT

In this study of 9212 eligible adult medical discharges from a teaching hospital, we developed and internally validated the HOSPITAL score that predicted the risk of 30day potentially avoidable readmission in medical patients with fair discriminatory power and good calibration. This easy-to-use model enables physicians to prospectively identify approximately 27% of the patients as highrisk of having a potentially avoidable readmission and would allow targeting intensive transitional care interventions to patients who might benefit the most.

The 22.3% overall readmission rate and 8.5% avoidable rate in our cohort are consistent with the findings of previous studies.^{1,2} As expected and shown in prior studies of all-cause or unplanned readmissions,^{8,9,17} the number of prior hospitalizations and the length of stay of the index admission were important predictors of po-

Table 3. HOSPITAL Score for 30-Day Potentially Avoidable Readmissions^a

Attribute	Points
Low <i>h</i> emoglobin level at discharge (<12 g/dL)	1
Discharge from an <i>o</i> ncology service	2
Low sodium level at discharge (<135 mEq/L)	1
Procedure during hospital stay (any <i>ICD-9-CM</i> coded procedure)	1
Index admission type: nonelective	1
No. of hospital <i>a</i> dmissions during the previous year	
0	0
1-5	2
>5	5
Length of stay \geq 5 d	2

Abbreviation: ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification.

SI conversion factors: To convert hemoglobin to grams per liter, multiply by 10; conversion of serum sodium to millimoles per liter is 1:1.

^aMaximum score, 13 points.

tentially avoidable readmission. A plausible explanation is that previous hospitalization may account for the total burden of illness, illness severity, functional status, and/or social environment. Length of stay also represents the severity of illness.

Even though all planned readmissions for chemotherapy were excluded from the outcome, being discharged from an oncology service was still associated with a high risk of potentially avoidable readmission. Patients discharged from an oncology division may be more fragile than most other medical service patients, and many readmissions may result from infectious complications, fluid and electrolyte abnormalities, a decrease in functional status, or end-of-life issues that might be avoidable. The Charlson Comorbidity Index or Elixhauser Comorbidity Index used in many prediction models is driven in large part by a diagnosis of cancer. Allaudeen et al¹⁰ and Silverstein et al¹⁴ found that the specific Elixhauser Comorbidity Index diagnosis of cancer was significantly associated with a 30-day readmission. It remains to be seen whether this finding is unique to large academic medical centers and/or institutions affiliated with a major cancer center, where the proportion of patients with cancer and the severity of disease are high.

Nonelective admissions seem to be followed by a higher risk for potentially avoidable readmission when compared with elective admissions. This may reflect the stability of the patient and is consistent with the findings of van Walraven et al.⁹

To our knowledge, no previous studies included the number of procedures performed during the index stay in their models. Depending on the procedure, this predictor may reflect disease severity as much as or more than avoidable complications of the procedure.

Finally, sodium and hemoglobin levels are markers of general prognosis (eg, in patients with heart failure, pulmonary embolism, and pneumonia).¹⁸⁻²⁰ To our knowledge, no other studies have included the sodium level as a predictor of readmission. Anemia was shown to be associated with hospital readmission within 90 days of discharge in one study.²¹

JAMA INTERN MED/VOL 173 (NO. 8), APR 22, 2013 WWW.JAMAINTERNALMED.COM 636

Table 4. Observed vs Predicted 30-Day Po	tentially Avoidable Readmissions
--	----------------------------------

Points		No. (%)			
	Risk Category	Patients in Each Category	Observed Proportion of Readmission	Estimated Risk of Readmission	
Derivation set (n = 6141)					
0-4	Low	3027 (49.3)	5.4	5.2	
5-6	Intermediate	1617 (26.3)	9.0	9.8	
≥7	High	1497 (24.4)	18.7	18.3	
Validation set (n = 3071)	·	× ,			
0-4	Low	1428 (46.5)	4.6	5.2	
5-6	Intermediate	855 (27.8)	9.7	9.8	
≥7	High	788 (26.7)	18.2	18.0	

Surprisingly, none of the most frequent comorbidities in these patients (based on index admission ICD-9-CM codes) was retained in the final model. Only congestive heart failure was significantly associated with a higher risk of 30-day potentially avoidable readmission, but this variable was not highly significant compared with most other covariates. This finding contrasts with some previous studies.^{7,10} It is possible that laboratory values, the number of previous admissions, the index length of stay, the need for inpatient procedures, nonelective admission, and oncology service are better predictors for the severity of illness and instability of the patient and reflect better the risk of readmission than a comorbidity measure does. The hypothesis that comorbidities or causes of admission do not matter as much as illness severity or clinical instability is attractive and has intuitive appeal. Another explanation could be the use of potentially avoidable readmissions as the outcome as opposed to the outcomes of previous studies.

Of the 26 unique prediction models identified in a recent systematic review,²² only 14 used the standard 30day outcome, which is more relevant for targeting interventions and for public reporting. Of these models, 6 are for patients with specific diseases, such as heart failure or pneumonia, which makes them difficult to apply, especially in patients with multiple conditions. Among the remaining 8 models, 6 include both medical and surgical patients, who may have different predictors of potentially avoidable readmissions. Also, of these 8 models, only 3 have a C statistic value greater than 0.65,^{7,9,11} and they might be good for risk adjustment but are less useful for prospectively targeting interventions because they do not focus on potentially avoidable readmissions and/or include predictors not available before discharge (eg, billing-based comorbidity scores). These limitations make these models less useful as a tool to target transitional care.

In comparison, the HOSPITAL score has fair discriminatory power and good calibration in identifying the risk of potentially avoidable readmission. It was developed using readmissions that might be prevented and are therefore potentially actionable. It can be used for all medical patients regardless of their main cause of admission. Finally, and most important, the HOSPITAL score is able to indicate the risk before a patient is discharged to allow targeting a timely transitional care intervention. We believe, of course, that all patients should receive highquality transitional care that meets certain standards. We are not recommending that low- and average-risk patients be deprived of effective transitional interventions. However, certain interventions that have been shown to be successful are resource intensive.^{3,4,23} One way to make best use of those limited resources is to reserve them for patients most likely to benefit.

These data must be interpreted in the context of this study's limitations. First, we were unable to identify readmissions that occurred outside our hospital network. However, previous studies^{24,25} using 30-day follow-up telephone calls and access to all available medical records have shown that more than 80% of Brigham and Women's Hospital medical patients are readmitted to 1 of the 3 hospitals in the Partners HealthCare network evaluated in this study. Moreover, there is no obvious reason to believe that the potentially avoidable readmissions outside the network might have such different predictors to change the current model.

Second, although we excluded inpatient deaths, postdischarge deaths were not identified or included in the outcome. Death within 30 days of discharge is a rare event (0.7% in one study),⁹ so a model predictive of potentially avoidable death or readmission would be unlikely to differ from a model predicting only potentially avoidable readmission.

Third, although SQLape has good performance characteristics in Switzerland, it has not been validated in the United States. We partially compensated for this by manually excluding planned readmissions (eg, because of coding errors). Other limitations in coding (eg, failure to document a complication of treatment) and in the algorithm itself mean that SQLape cannot perfectly distinguish avoidable from unavoidable readmissions, but it did allow us to create a cohort enriched with avoidable readmissions and develop a model better able to identify predictors of avoidable readmission than would otherwise be the case.

Fourth, it is likely that our model excluded important predictors of potentially avoidable readmission, such as functional status, health literacy, degree of social support, and previous medication adherence.⁷ However, we chose not to include these predictors, which are infrequently measured and often difficult to obtain, because the study's goal was to derive a model that could be easily and widely used.

Fifth, hemoglobin and sodium values are available only near the time of discharge and might limit the time available to effectively deploy an intensive discharge intervention. However, we calculated that 60% of patients defined as high risk could be identified 2 or more days before discharge, based on information available at that time, allowing sufficient time for intervention deployment. Similarly, procedures were technically defined by postdischarge *ICD-9-CM* codes (Table 2), but these could be recognized easily by clinicians or administrators during the hospitalization as any procedure requiring patient consent.

Sixth, this study performed only an internal validation of the model. An external validation of the HOSPITAL score is likely required, particularly in smaller hospitals, before widespread implementation.

Last, predicting potentially avoidable readmissions is only a proxy for identifying who might benefit from specific interventions. Intervention studies targeting this patient population need to be done to definitively prove its usefulness.

In conclusion, we propose a prediction model, HOSPITAL, that provides a practical tool to assess 30day potentially avoidable readmission risk in medical patients. The use of this simple score before discharge may help target transitional care for patients who might benefit the most and consequently reduce the rate of avoidable readmission.

Accepted for Publication: October 26, 2012.

Published Online: March 25, 2013. doi:10.1001/jamainternmed.2013.3023

Correspondence: Jacques Donzé, MD, MSc, Division of General Medicine and Primary Care, Brigham and Women's Hospital, 1620 Tremont St, OBC-3-2U, Boston, MA 02120 (jdonze@partners.org).

Author Contributions: Study concept and design: Donzé and Schnipper. Acquisition of data: Donzé and Williams. Analysis and interpretation of data: All authors. Drafting of the manuscript: Donzé. Critical revision of the manuscript for important intellectual content: All authors. Statistical analysis: Donzé. Obtained funding: Donzé. Administrative, technical, and material support: Williams. Study supervision: Aujesky and Schnipper.

Conflict of Interest Disclosures: Dr Schnipper is a consultant to QuantiaMD, for whom he has helped create online educational materials for providers and patients regarding patient safety, including medication safety during transitions in care. The findings of this study are not a part of those materials. Dr Schnipper has received grant funding from sanofi-aventis for an investigator-initiated study to design and evaluate an intensive discharge and follow-up intervention for patients with diabetes mellitus. **Funding/Support:** Dr Donzé was supported by grants PBLAP3-131814 and PBLAP3-136815 from the Swiss Science National Foundation and by the SICPA Foundation. **Role of the Sponsors:** The Swiss Science National Foundation and the SICPA Foundation had no role in the design and conduct of this study, the analysis or interpreta-

tion of the data, or the preparation of the manuscript. **Previous Presentation**: A portion of the study's findings were presented at the Annual Meeting of the Society of General Internal Medicine; May 12, 2012; Orlando, Florida.

Online-Only Material: The eTables are available at http: //www.jamainternalmed.com. Additional Contributions: Yves Eggli, MD, from the Institute of Health Economics and Management of the University of Lausanne, Switzerland, screened the database for potentially avoidable readmissions using the SQLape algorithm.

REFERENCES

- Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. N Engl J Med. 2009;360(14):1418-1428.
- van Walraven C, Bennett C, Jennings A, Austin PC, Forster AJ. Proportion of hospital readmissions deemed avoidable: a systematic review. *CMAJ*. 2011;183 (7):E391-E402.
- Jack BW, Chetty VK, Anthony D, et al. A reengineered hospital discharge program to decrease rehospitalization: a randomized trial. *Ann Intern Med.* 2009; 150(3):178-187.
- Naylor MD, Brooten D, Campbell R, et al. Comprehensive discharge planning and home follow-up of hospitalized elders: a randomized clinical trial. JAMA. 1999; 281(7):613-620.
- tj;2Coleman EA, Parry C, Chalmers S, Min SJ. The care transitions intervention: results of a randomized controlled trial. Arch Intern Med. 2006;166(17):1822-1828.
- Kane RL, Keckhafer G, Flood S, Bershadsky B, Siadaty MS. The effect of Evercare on hospital use. J Am Geriatr Soc. 2003;51(10):1427-1434.
- Coleman EA, Min SJ, Chomiak A, Kramer AM. Posthospital care transitions: patterns, complications, and risk identification. *Health Serv Res.* 2004;39(5):1449-1465.
- Hasan O, Meltzer DO, Shaykevich SA, et al. Hospital readmission in general medicine patients: a prediction model. J Gen Intern Med. 2010;25(3):211-219.
- van Walraven C, Dhalla IA, Bell C, et al. Derivation and validation of an index to predict early death or unplanned readmission after discharge from hospital to the community. CMAJ. 2010;182(6):551-557.
- Allaudeen N, Vidyarthi A, Maselli J, Auerbach A. Redefining readmission risk factors for general medicine patients. J Hosp Med. 2011;6(2):54-60.
- Halfon P, Eggli Y, van Melle G, Chevalier J, Wasserfallen JB, Burnand B. Measuring potentially avoidable hospital readmissions. *J Clin Epidemiol.* 2002; 55(6):573-587.
- Halfon P, Eggli Y, Prêtre-Rohrbach I, Meylan D, Marazzi A, Burnand B. Validation of the potentially avoidable hospital readmission rate as a routine indicator of the quality of hospital care. *Med Care.* 2006;44(11):972-981.
- Boult C, Dowd B, McCaffrey D, Boult L, Hernandez R, Krulewitch H. Screening elders for risk of hospital admission. J Am Geriatr Soc. 1993;41(8):811-817.
- Silverstein MD, Qin H, Mercer SQ, Fong J, Haydar Z. Risk factors for 30-day hospital readmission in patients ≥65 years of age. *Proc (Bayl Univ Med Cent)*. 2008; 21(4):363-372.
- Moons KG, Harrell FE, Steyerberg EW. Should scoring rules be based on odds ratios or regression coefficients? *J Clin Epidemiol.* 2002;55(10):1054-1055.
- Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology*. 1982;143(1):29-36.
- Billings J, Dixon J, Mijanovich T, Wennberg D. Case finding for patients at risk of readmission to hospital: development of algorithm to identify high risk patients. *BMJ*. 2006;333(7563):327. doi.org/10.1136/bmj.38870.657917.AE.
- Groenveld HF, Januzzi JL, Damman K, et al. Anemia and mortality in heart failure patients: a systematic review and meta-analysis. *J Am Coll Cardiol.* 2008; 52(10):818-827.
- Donzé J, Labarère J, Méan M, Jiménez D, Aujesky D. Prognostic importance of anaemia in patients with acute pulmonary embolism. *Thromb Haemost*. 2011; 106(2):289-295.
- Waterer GW, Kessler LA, Wunderink RG. Medium-term survival after hospitalization with community-acquired pneumonia. *Am J Respir Crit Care Med.* 2004; 169(8):910-914.
- Smith DM, Norton JA, McDonald CJ. Nonelective readmissions of medical patients. J Chronic Dis. 1985;38(3):213-224.
- Kansagara D, Englander H, Salanitro A, et al. Risk prediction models for hospital readmission: a systematic review. JAMA. 2011;306(15):1688-1698.
- Naylor MD, Brooten DA, Campbell RL, Maislin G, McCauley KM, Schwartz JS. Transitional care of older adults hospitalized with heart failure: a randomized, controlled trial. *J Am Geriatr Soc.* 2004;52(5):675-684.
- Schnipper JL, Roumie CL, Cawthon C, et al; PILL-CVD Study Group. Rationale and design of the Pharmacist Intervention for Low Literacy in Cardiovascular Disease (PILL-CVD) study. *Circ Cardiovasc Qual Outcomes*. 2010;3(2):212-219.
- Kripalani S, Roumie CL, Dalal AK, et al; PILL-CVD (Pharmacist Intervention for Low Literacy in Cardiovascular Disease) Study Group. Effect of a pharmacist intervention on clinically important medication errors after hospital discharge: a randomized trial. *Ann Intern Med.* 2012;157(1):1-10.

JAMA INTERN MED/VOL 173 (NO. 8), APR 22, 2013 WWW.JAMAINTERNALMED.COM 638