

**Correspondence:** Dr O'Malley, Department of Medicine, Division of General Internal Medicine, Walter Reed Army Medical Center, 6900 Georgia Ave, Washington, DC 20307 (pomalley@usuhs.mil).

**Financial Disclosure:** None reported.

**Disclaimer:** The views expressed herein are those of the author only and are not to be construed as those of the Department of the Army or Department of Defense.

**Additional Contributions:** Mitchell Katz, MD, provided helpful edits and comments.

## REFERENCES

1. Katon WJ, Lin EH, Von Korff M, et al. Collaborative care for patients with depression and chronic illnesses. *N Engl J Med*. 2010;363(27):2611-2620.
2. Chisholm-Burns MA, Kim Lee J, Spivey CA, et al. US pharmacists' effect as team members on patient care: systematic review and meta-analyses. *Med Care*. 2010; 48(10):923-933.
3. Callahan CM, Boustani MA, Unverzagt FW, et al. Effectiveness of collaborative care for older adults with Alzheimer disease in primary care: a randomized controlled trial. *JAMA*. 2006;295(18):2148-2157.
4. GESICA Investigators. Randomised trial of telephone intervention in chronic heart failure: DIAL trial. *BMJ*. 2005;331(7514):425.
5. Dobscha SK, Corson K, Perrin NA, et al. Collaborative care for chronic pain in primary care: a cluster randomized trial. *JAMA*. 2009;301(12):1242-1252.
6. Rollman BL, Belnap BH, Mazumdar S, et al. A randomized trial to improve the quality of treatment for panic and generalized anxiety disorders in primary care. *Arch Gen Psychiatry*. 2005;62(12):1332-1341.
7. DeBusk RF, Miller NH, Parker KM, et al. Care management for low-risk patients with heart failure: a randomized, controlled trial. *Ann Intern Med*. 2004;141(8):606-613.
8. Laramee AS, Levinsky SK, Sargent J, Ross R, Callas P. Case management in a heterogeneous congestive heart failure population: a randomized controlled trial. *Arch Intern Med*. 2003;163(7):809-817.
9. Krein SL, Klamerus ML, Vijan S, et al. Case management for patients with poorly controlled diabetes: a randomized trial. *Am J Med*. 2004;116(11):732-739.
10. Dobscha SK, Corson K, Hickam DH, Perrin NA, Kraemer DF, Gerrity MS. Depression decision support in primary care: a cluster randomized trial. *Ann Intern Med*. 2006;145(7):477-487.
11. Fihn SD, Bucher JB, McDonell M, et al. Collaborative care intervention for stable ischemic heart disease. *Arch Intern Med*. 2011;171(16):1471-1479.
12. Pape GA, Hunt JS, Butler KL, et al. Team-based care approach to cholesterol management in diabetes mellitus: 2-year cluster randomized controlled trial. *Arch Intern Med*. 2011;171(16):1480-1486.

## Correction

**Typographical Error in Abstract.** In the "Background" section of the abstract of the Original Investigation "Impact of QRS Duration on Clinical Event Reduction With Cardiac Resynchronization Therapy: Meta-analysis of Randomized Controlled Trials" by Sipahi et al, published in the September 12, 2011, issue of the *Archives* (2011;171[16]:1454-1462) and published online June 13, 2011, the reported range for moderately prolonged QRS duration was incorrect. The correct range is 120 to 149 milliseconds.